PB81-192197

# Environmental Assessment of Buccaneer Gas and Oil Field in the Northwestern Gulf of Mexico 1975-1980. Volume VI. Trace Metals

(U.S.) National Marine Fisheries Service
Galveston, TX

Prepared for

Environmental Protection Agency
Washington, DC

Nov 80

U.S. Department of Commerce
National Technical Information Service
NTIS

PB81-192197

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA 22161

NOAA FORM 25-13 (1-78)

**BIBLIOGRAPHIC DATA SHEET**

U. S. DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION

| 1. NOAA ACCESSION NUMBER | 2. | 3. RECIPIENT'S ACCESSION NUMBER |
|---|---|---|
| NOAA-81012802 | | PB81 192197 |

**4. TITLE AND SUBTITLE**
Environmental Assessment of Buccaneer Gas and Oil Field in the Northwestern Gulf of Mexico, 1975-1980. Volume VI: Trace Metals

**5. REPORT DATE**
Nov 1980

**6.**

**7. AUTHOR(S)**
J.B. Tillery (Southwest Res. Ins., San Antonio, TX)
William B. Jackson and Wilkens, E. Peter (Eds.)

**8. REPORT NO.**
NOAA-TM-NMFS-SEFC-52

**9. PERFORMING ORGANIZATION NAME AND ADDRESS**
NOAA, National Marine Fisheries Service,
Galveston, TX 77550,
Southeast Fisheries Center

**10. PROJECT/TASK NO.**

**11. CONTRACT/GRANT NO.**
Interagency Agreement
EPA-1AG-D5-E693-EO

**12. SPONSORING ORGANIZATION NAME AND ADDRESS**
Same; and
U.S. Environmental Protection Agency

**13. TYPE OF REPORT AND PERIOD COVERED**

**14.**

**15. PUBLICATION REFERENCE**
NOAA Technical Memorandum NMFS-SEFC-52, November 1980. 70 p, numerous fig, tab, ref.

**16. ABSTRACT** This report provides an overview of four years (1976-1980) of trace metal investigations conducted as a part of a larger multidisciplinary, marine, environmental assessment program of the Buccaneer gas and oil field. Concentrations of Ag, Al, As, Ba, Be, Cd, Co, Cr, Cu, Hg, Fe, Mn, Ni, Pb, Sb, Se, Sr, Tl and Zn have been determined in surficial sediments, subsurface sediments, suspended particulate matter, seawater, produced brine, crude oil and various tissues of biological organisms collected seasonally near two petroleum production platforms. Concentrations of Ba, Cd, Cr, Co, Cu, Mn, Pb, Sr, Hg and Zn in surficial sediments have been related to the platforms or activities on them. Trace metal concentrations in suspended particulate matter are higher than in bottom sediments. Seawater trace metal concentrations are within the range reported for shelf waters. Produced brine discharge have concentrations of Ba, Cd, Cr, Fe, Hg, Mn, Sr, Tl and Zn that are higher than seawater and vary with time. No evidence of excessive bioaccumulation of trace metals in marine organisms from the area around the production platforms was established. Seasonal variations of trace metals were observed in various marine organisms. (Author)

**17. KEY WORDS AND DOCUMENT ANALYSIS**

**17A. DESCRIPTORS**

*Metals, *Environmental effects, Sediments, Gulf of Mexico, Oil fields, Biota, Gases, Water pollution effects

**17B. IDENTIFIERS/OPEN-ENDED TERMS**

Hydrocarbons, Gas fields, Trace metals

**17C. COSATI FIELD/GROUP**

6F, 8A COWRR 05C

**18. AVAILABILITY STATEMENT**
Released for distribution:
Elaine L Downs

| 19. SECURITY CLASS (This report) | 21. NO. OF PAGES |
|---|---|
| UNCLASSIFIED | 71 p. |
| **20. SECURITY CLASS (This report)** | **22. PRICE** |
| UNCLASSIFIED | |

NOAA FORM 25-13 (1-78) SUPERSEDES ALL PREVIOUS EDITIONS.

# NOAA Technical Memorandum NMFS-SEFC-52

## Environmental Assessment of Buccaneer Gas and Oil Field In the Northwestern Gulf of Mexico, 1975-1980.

## VOL. VI - TRACE METALS


BY

J.B. Tillery

Southwest Research Institute

6228 Culebra

San Antonio, Texas 78284



A report to the Environmental Protection Agency on work conducted under provisions of Interagency Agreement EPA-IAG-D5-E693-EO during 1975-1980.



EDITORS

William B. Jackson

Senior Advisor

Contracts & Deliverables

and

E. Peter Wilkens

Fishery Biologist



U. S. DEPARTMENT OF COMMERCE

Philip M. Klutznick, Secretary

National Oceanic and Atmospheric Administration

Richard A. Frank, Administrator

National Marine Fisheries Service

Terry L. Leitzell, Assisistant Administrator for Fisheries



This TM series is used for documentation and timely communication of preliminary results, interim reports, or similar special purpose information. Although the memos are not subject to complete formal review, editorial control, or detailed editing, they are expected to reflect sound professional work.

## DISCLAIMER

This document is a Milestone Report. It has been reviewed by the National Marine Fisheries Service and the National Oceanic and Atmospheric Administration and approved for printing. Such approval does not signify that the contents necessarily reflect the views and policies of the U. S. Environmental Protection Agency, NOAA or NMFS. This Report has not been formally released by the EPA. Mention of trade names or commerical products herein does not constitute endorsement or recommendation for use.

NOTICE

This document is a Milestone Report. It has not been formally released by the U.S. Environmental Protection Agency and should not at this stage be construed to represent Agency policy.

This volume should be cited as follows:

Tillery, J. B. 1980. Trace metals. Vol. VI. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Technical Memorandum NMFS-SEFC-52. 39 p. Available from: NTIS, Springfield, Virginia.

Volume VI - TRACE METALS

TABLE OF CONTENTS


I. Editors' Section

| | Page |
|---|---|
| Project Administration | v |
| List of Volumes | vii |
| Foreword | ix |
| List of Reports and Publications | xi |
| Published Reports | xi |
| Dissertations and Theses | xvi |
| Publications in Press or in Preparation | xvi |
| List of Archived Data | xviii |
| Introduction | xxi |
| Location of Study Area | xxi |
| Operation History of Buccaneer Field | xxi |
| Fig. 1. Location of Buccaneer Field | xxii |
| Fig. 2. Buccaneer Field Structures | xxiv |
| Fig. 3. Shell Oil Company's Alphanumerical Identification of Buccaneer Field Structures | xxv |

II. Principal Investigators' Section

| | |
|---|---|
| Work Unit 2.4.2 Trace Metals | xxvi |

## PROJECT ADMINISTRATION

NOAA

Program Manager

W. Lawrence Pugh
Oceans Program Office
Rockville, Maryland

NMFS

Contracting Officer's Technical Representative

Edward F. Klima, Ph.D.
Director
Galveston Laboratory
Southeast Fisheries Center

Project Manager

Charles W. Caillouet, Ph.D.
Chief, Environmental Research Division

Project Staff (Environmental Research Division)

William B. Jackson
Senior Advisor
Contracts and Deliverables

Gregg R. Gitschlag
Senior Advisor
Field Operations and Logistics

E. Peter H. Wilkens
Fishery Biologist

Gary M. Faw
Fishery Biologist

Robert M. Avent, Ph.D.
Oceanographer

Dennis Koi
Computer Programmer

Petronila C. Prado
Clerk Stenographer

Mary Taylor
Clerk Typist

Patsy Hunter
Clerk Typist

Susan Gray
Clerk Typist

Beatrice Richardson
Clerk Typist

Leesa Young
Biological Aide

Julie Mellen
Student Aide

Richard Devereux
Coop. Student Biologist

## LIST OF VOLUMES

This Milestone Report is printed in six separate volumes:

Volume I - SEDIMENTS, PARTICULATES AND VOLATILE HYDROCARBONS

Work Unit 2.3.2 — Investigations of Surficial Sediments, Suspended Particulates and Volatile Hydrocarbons at Buccaneer Gas and Oil Field

Texas A&M University

J. Brooks, Ph.D.
E. Estes, Ph.D.
D. Wiesenburg
C. Schwab
H. Abdel-Reheim

Volume II - FISHES AND MACRO-CRUSTACEANS

Work Unit 2.3.5/ 2.3.8 — Pelagic, Reef and Demersal Fishes, and Macro-crustaceans/Biofouling Communities

LGL Ecological Research Associates, Inc.

B. Gallaway, Ph.D.

Volume III - BACTERIA

Work Unit 2.3.7 — Bacteriology of a Gulf of Mexico Gas and Oil Field

University of Houston

R. Sizemore, Ph.D.
K. Olsen

Volume IV - CURRENTS AND HYDROGRAPHY

Work Unit 2.3.9 — Currents Patterns and Hydrography of the Buccaneer Field and Adjacent Waters

NMFS Atlantic Environmental Group

R. Armstrong

Volume V - HYDROCARBONS

Work Unit 2.4.1 — Hydrocarbons, Biocides and Sulfur

University of Houston

B. Middleditch, Ph.D.

Volume VI - TRACE METALS

Work Unit 2.4.2 — Trace Metals

Southwest Research Institute

J. Tillery

# FOREWORD

Increased petroleum development of the outer continental shelf (OCS) of the United States is anticipated as the U.S. attempts to reduce its dependency on foreign petroleum supplies. To obtain information concerning the environmental consequences of such development, the Federal Government has supported major research efforts on the OCS to document environmental conditions before, during, and after oil and gas exploration, production, and transmission. Among these efforts is the Environmental Assessment of Buccaneer Gas and Oil Field in the Northwestern Gulf of Mexico, a project funded by the Environmental Protection Agency (EPA) through interagency agreement with the National Oceanic and Atomospheric Administration (NOAA) and managed by the National Marine Fisheries Service (NMFS), Southeast Fisheries Center (SEFC), Galveston Laboratory, in Galveston, Texas. Initiated in the autumn of 1975, the study was completed in 1980. Its major products have been annual reports disseminated by the National Technical Information Service, data files archived and disseminated by NOAA's Environmental Data and Information Service, and research papers written by participating investigators and published in scientific or technical journals. Results have also been made available through EPA/NOAA/NMFS project reviews and workshops attended by project participants, and various governmental (Federal and State), private, and public user groups. The final product are these milestone reports summarizing the findings of the major investigative components of the study.

Objectives of the project were (1) to identify and document the types and extent of biological, chemical and physical alterations of the marine ecosystem associated with Buccaneer Gas and Oil Field, (2) to determine specific pollutants, their quantity and effects, and (3) to develop the capability to describe and predict fate and effects of Buccaneer Gas and Oil Field contaminants. The project used historical and new data and included investigations both in the field and in the laboratory. A brief Pilot Study was conducted in the autumn and winter of 1975-76, followed by an extensive biological/chemical/physical survey in 1976-77 comparing the Buccaneer Gas and Oil Field area with adjacent undeveloped or control areas. In 1977-78, investigations were intensified within Buccaneer Gas and Oil Field, comparing conditions around production platforms, which release various effluents including produced brine, with those around satellite structures (well jackets) which release no effluents. In 1978-79, studies around Buccaneer Gas and Oil Field structures focused on (1) concentrations and effects of pollutants in major components of

the marine ecosystem, including seawater, surficial sediments, suspended particulate matter, fouling community, bacterial community, and fishes and macro-crustaceans, (2) effects of circulation dynamics and hydrography on distribution of pollutants, and (3) mathematical modeling to describe and predict sources, fate and effects of pollutants. The final year, 1979-80, of study continued to focus on items (1) and (2) and on preparation of the milestone reports which represented the final products of this study.

This project has provided a unique opportunity for a multi-year investigation of effects of chronic, low-level contamination of a marine ecosystem associated with gas and oil production in a long-established field. In many respects, it represents a pioneering effort. It has been made possible through the cooporation of government agencies, Shell Oil Company (which owns and operates the field) and various contractors including universities and private companies. It is anticipated that the results of this project will impact in a significant way on future decisions regarding operations of gas and oil fields on the OCS.

Charles W. Caillouet, Project Manager
Chief, Environmental Research Division
and
William B. Jackson and E. Peter Wilkens,
Editors

## LIST OF REPORTS AND PUBLICATIONS

### Published Reports


Armstrong, R. S. 1980. Current patterns and hydrography . Vol. IV. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental Assessment of Buccaneer gas and oilfield in the northwestern Gulf of Mexico, 1975-1980. NOAA Technical Memorandum NMFS-SEFC-50. p. Available from: NTIS, Springfield, Virginia.

Brooks, J. M., E. L. Estes, D. A. Wiesenburg , C. R. Schwab and H. A. Abdel-Reheim. 1980. Investigations of surficial sediments, suspended particulates and volatile hydrocarbons at Buccaneer gas and oil field. Vol I. In: Jackson, W. B. and E. P. Wilkens, (eds.). Environmental Asessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Technical Memorandum NMFS-SEFC-47, 89 p. Available from: NTIS, Springfield Virginia.

Brooks, J. M. , E. L. Estes and W. Huang. 1980. Investigations of surficial sediments and suspended particulates at Buccaneer field. Vol. II. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-36, 261 p. Available from: NTIS, Springfield, Virginia.

Danek, L. J. and M. S. Tomlinson. 1980. Currents and hydrography of the Buccaneer field and adjacent waters. Vol. VI. In: Jackson, W. B. and E. P. Wilkens, (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-40, 33 p. Available from: NTIS, Springfield, Virginia.

Finucane, J. H. 1977. Ichthyoplankton of the Buccaneer oil field off Galveston, Texas. Paper presented at the Fourth Biennial International Estuarine Research Conference, Mt. Pocono, Pa. (abstract published in proceedings).

Fucik, K. and I. T. Show, Jr. 1980. Sources, fate and effects modeling. Vol. IX. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-43, p. Available from: NTIS, Springfield, Virginia.

Gallaway, B. J. 1980. Pelagic, reef and demersal fishes and macrocrustaceans/biofouling communities. Vol II. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental Assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Technical Memorandum NMFS-SEFC-48, 82 p. Available from: NTIS, Springfield, Virginia.

Gallaway, B. J., R. Howard, K. Green and L. Martin. 1976. A study plan for the Buccaneer oil field assessment program. A report to NMFS from LGL Limited-U.S., Inc., Bryan, Texas, 69 pp.

Gallaway, B. J. and L. R. Martin. 1980. Effect of gas and oil field structures and effluents on pelagic and reef fishes, and demersal fishes and macrocrustaceans. Vol. III. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-37, 49 p. Available from: NTIS, Springfield, Virginia.

Harper, D. E., Jr., R. J. Scrudato and C. S. Giam. 1976. A preliminary environmental assessment of the Buccaneer oil/gas field (pilot study of the Buccaneer oil field (benthos and sediments). A final report from Texas A&M University to the National Marine Fisheries Service, Galveston, Texas (Contract No. 03-6-042-35110), 63 pp.

Hollaway, S. J., G. M. Faw, R. K. Sizemore. 1980. The bacterial community composition of an active oil field in the northwestern Gulf of Mexico. Mar. Poll. Bull. 11:153-156.

Howard, R. L, G. S. Boland, B. J. Gallaway and G. D. Dennis. 1980. Effects of gas and oil field structures and effluents on fouling community production and function. Vol. V. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-39, 60 p. Available from: NTIS, Springfield, Virginia.

Jackson, W. B. (Editor). 1977. Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1976-1977. NOAA Annual Report to EPA, Project Number EPA-IAG-D5-E693-EO, 759 pp. Available from: NTIS, Springfield, Virginia, Accession No. PB283890.

Jackson, W. B. (Editor). 1979. Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Volume I: Synopsis. NOAA Annual Report to EPA, Project Number

EPA-IAG-D5-E693-EO. 62 pp. Available from: NTIS, Springfield, Virginia, Accession No. PB301339/AS.

Jackson, W. B. (Editor). 1979. Environmental assessement of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Volume III: Chemical and physical investigations. NOAA Annual Report to EPA, Project Number EPA-IAG-D5-E693-EO. 722 pp. Available from: NTIS, Springfield, Virginia, Accession No. PB80107899.

Jackson, W. B. (Editor). 1979. Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977-1978. Volume II: Data management and biological investigations. NOAA Annual Report to EPA, Project Number EPA-IAG-D5-E693-EO, 799 pp. Available from: NTIS, Springfield, Virginia, Accession No. PB80165970.

Jackson W. B. and E. P. Wilkens. (Editors). 1980. Synopsis. Vol. I. In: Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-35, p. Available from: NTIS, Springfield, Virginia.

Jackson, W. B., K. N. Baxter and C. W. Caillouet. 1978. Eviron-mental assessment of the Buccaneer oil and gas field off Galveston, Texas: An overview. Proceedings of the Offshore Technology Conference, May, 1978, Houston, Texas, OTC 3081:277-284.

Jackson, W. B., K. N. Baxter, and C. W. Caillouet. 1979. Environ-mental assessment of an offshore oil field. Ecolibrium 8(1): 7-9.

McCall, J. O. and R. K. Sizemore. 1979. Description of a bac-teriocinogenic plasmid in Beneckea harveyi. Appl. Environ. Microbiol. 38(5):974-979.

McKinney, L. 1979. Liljeborgiid amphiopods from the Gulf of Mexico and Caribbean Sea. Bull. Mar. Sci. 29(2):140-154.

Middleditch, B. S. 1980. Hydrocarbons, biocides, and sulfur. Vol. III. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Technical Memorandum NMFS-SEFC-51, 70 p. Available from NTIS, Springfield, Virginia.

Middleditch, B. S. and B. Basile. 1976. Deuteriated analogs as internal standards for the quantitation of environmental alkanes by gas chromatography. Analyt. Letters 9:1031-1034.

Middleditch, B. S. and B. Basile. 1979. Sulfur-34 as an internal standard for quantitation of environmental sulfur by combined gas chromatography-mass spectometry. Analyt. Letters 12(A):777-781.

Middleditch, B. S., B. Basile and E. S. Chang. 1977. Environmental effects of offshore oil production: alkanes in the region of the Buccaneer oil field. J. Chromatography 142(1977):777-785.

Middleditch, B. S., B. Basile and E. S. Chang. 1978. Alkanes in surficial sediments from the region of the Buccaneer oil field. J. Chromatography 158:449-463.

Middleditch, B. S., B. Basile and E. S. Chang. 1978. Discharge of alkanes during offshore oil production in the Buccaneer oil field. Bull. Environ. Contam. Toxicol. 20:59-65.

Middleditch, B. S., B. Basile and E. S. Chang. 1979. Alkanes in seawater in the vicinity of the Buccaneer oil field. Bull. Environ. Contam. Toxicol. 21:413-420.

Middleditch, B. S., E. S. Chang and B. Basile. 1979. Alkanes in barnacles (Balanus tintinnabulum) from the Buccaneer oil field. Bull. Environ. Contam. Toxicol. 24:6-12.

Middleditch, B. S., E. S. Chang and B. Basile. 1979. Alkanes in plankton from the Buccaneer oil field. Bull. Environ. Contam. Toxicol. 21:421-427.

Middleditch, B. S., E. S. Chang, B. Basile, and S. R. Missler. 1979. Alkanes in fish from the Buccaneer oil field. Bull. Environ. Contam. Toxicol. 22:249-257.

Middleditch, B. S. and D. West. 1979. Bicyclohexyl interference in the quantitation of environmental alkanes by gas chromatography. Analyt. Letters 12:1279-1282.

Middleditch, B. S. and D. West. 1980. Hydrocarbons, biocides, and sulfur. Vol. VII. In: Jackson, W. B. and E. P. Wilkens, (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-41, p. Available from: NTIS, Springfield, Virginia.

National Marine Fisheries Service. 1977. Environmental assessment of an active oil field in the northwestern Gulf of Mexico. A report of NMFS/LGL Workshop I, 1977-1978. NMFS Galveston Laboratory, Southeast Fisheries Center, NOAA, USDOC. Project Number EPA-IAG-D5-E693-EO.

Savastano, K. and H. Holley. 1980. Implement, monitor, and modify data management system. Vol. I. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental asessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-35. p. Available from: NTIS, Springfield, Virginia.

Show, I. T., Jr. 1979. An application of compartmental models to mesoscale marine ecosystems, pages 73-79. In: J. H. Matis, B. C. Patten, and G. C. White (eds.). Compartmental analysis of ecosystem models. International Co-operative Publishing House, Fairland, Maryland, U.S.A.

Sizemore, R. K. and K. Olsen. 1980. Bacterial communities. Vol. IV. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-38, 32 p. Available from: NTIS, Springfield, Virginia.

Sizemore, R. K. and K. Olsen. 1980. Bacteriology of a Gulf of Mexico gas and oil field. Vol. III. In: Jackson, W. B. and E. P. Wilkens (Eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Technical Memorandum NMFS-SEFC-49, 21 p. Available from: NTIS, Springfield, Virginia.

Smedes, G., J. Calman and J. Beebe. 1980. Hydrodynamic modeling. Vol. X. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-44, 57 p. Available from: NTIS, Springfield, Virginia.

Tillery, J. B. 1980. Trace metals. Vol. VIII. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1978-1979. NOAA Technical Memorandum NMFS-SEFC-42, 93 p. Available from: NTIS, Springfield, Virginia.

Tillery, J. B. 1980. Trace metals. Vol. VI. In: Jackson, W. B. and E. P. Wilkens (eds.). Environmental assessment of Buccaneer gas and oil field in the northwestern Gulf of Mexico, 1975-1980. NOAA Technical Memorandum NMFS-SEFC-52, 39 p. Available from: NTIS, Springfield, Virginia.

Wheeler, R. B., R. R. Schwarzer and J. B. Anderson. 1978. Assessment of environmental impact of offshore production in the Buccaneer oil field: sedimentologic and geochemical results. Proceedings

of the Offshore Technology Conference, May 1978, Houston, Texas OTC 3082:285-290.

## Dissertations and Theses

Basile, B. 1978. Environmental effects of offshore oil production. Ph.D. Dissertation, University of Houston, Houston, Texas.

Boland, G. S. 1980. Morphological parameters of the barnacle, Balanus tintinnabulum antillensis, as indicators of physiological and environmental conditions. M. S. Thesis, Texas A&M University, College Station, Texas.

Chang, E. S. 1978. Sterols in the marine food web. M.S. Thesis, University of Houston, Houston, Texas.

McCall, J. O. 1979. Isolation and characterization of a bacteriocinogenic strain of the bioluminescent marine bacterium Beneckea harveyi. M.S. Thesis, University of Houston, Houston, Texas.

West, D. L. 1979. Hydrocarbons, sulfur and acrolein in the marine ecosystem associated with offshore oil production. M.S. Thesis, University of Houston, Houston, Texas.

Wheeler, R. B. 1979. Environmental trace metal geochemistry of the Buccaneer oil and gas field. M.S. Thesis, Rice University, Houston, Texas. 176 pp.

## Publications in Press or in Preparation

Anderson, J. B., R. R. Schwarzer, B. S. Middleditch, and B. Basile. Trace metal and hydrocarbon distribution in the region of the Buccaneer oil field. M.S. in preparation.

Devereux, R. and R. K. Sizemore. Incidence of degradative plasmids in hydrocarbon-utilizing bacteria. M.S. submitted to Developements in Industrial Microbiology.

Hada, H. S. and R. K. Sizemore. A survey of the incidence of plasmids in marine Vibrio spp. M.S. submitted to Appl. Environ. Microbiol.

McKinney, L. 1980. The genus Photis from the Texas coast with a description of a new species, Photis melanicus. Contrib. Mar. Sci. (in press).

Middleditch, B. S. and B. Basile. 1980. Alkanes in shrimp from the Buccaneer oil field. Bull. Environ. Contam. Toxicol. (in press).

Middleditch, B. S. and B. Basile. 1980. Alkanes in benthic organisms from the Buccaneer oil field. Bull. Environ. Contam. Toxicol. (in press).

Middleditch, B. S. and B. Basile. 1980. Discharge of elemental sulfur and its distribution in the region of the Buccaneer oil field. J. Chromatog. (in press).

Olsen, K. D. and R. K. Sizemore. Effects of an established offshore oil platform on the autochthonous bacteria community. M. S. submitted to Developments in Industrial Microbiology.

Rose, C. D. and T. J. Ward. Principles of aquatic hazard evaluation as applied to ocean-disposed wastes. M.S. in preparation; intended for publication as a report for Amer. Soc. Testing. Materials.

Wheeler, R. B. 1980. Environmental trace metal geochemistry of the Buccaneer oil and gas field. Environ. Geol. (in press).

## LIST OF ARCHIVED DATA

Data available from U.S. Department of Commerce, NOAA, EDIS,
National Oceanographic Data Center, Washington, D.C. 20235

| Year | Data Type | NODC Accession Number |
|---|---|---|
| 1976-1977 | Demersal Fish | 78-0501 |
| 1976-1977 | Sediment | 78-0501 |
| 1976-1977 | Birds | 78-0501 |
| 1976-1977 | Ichthyoplankton | 78-0501 |
| 1976-1977 | Pelagic Fish | 78-0501 |
| 1976-1977 | Plankton | 78-0501 |
| 1976-1977 | Sessile Fauna | 78-0501 |
| 1976-1977 | Total Organics | 78-0501 |
| 1976-1977 | Hydrocarbons | 78-0501 |
| 1976-1977 | Fish Determination | 78-0501 |
| 1976-1977 | Ocean Serial Stations | 78-0501 |
| 1976-1977 | Trace Metals | 78-0501 |
| 1976-1977 | Benthos | 78-0501 |
| 1976-1977 | Drift Bottle Releases | 78-0501 |

| Year | Data Type | NODC Accession Number |
|---|---|---|
| 1977-1978 | Brine Dye Release | 80-0423 |
| 1977-1978 | Fish Bioassay | 80-0423 |
| 1977-1978 | Ichthyoplankton | 80-0423 |
| 1977-1978 | Food Habits-Station | 80-0423 |
| 1977-1978 | Food Habits-Stomach | 80-0423 |
| 1977-1978 | Reef Fish Census | 80-0423 |
| 1977-1978 | Pelagic Fish Census | 80-0423 |
| 1977-1978 | Biofouling | 80-0423 |
| 1977-1978 | Transponding Buoy (2 Files) | 80-0423 |
| 1977-1978 | Drift Bottle Release/Recovery | 80-0423 |
| 1977-1978 | Dye Study-station | 80-0423 |
| 1977-1978 | Ocean Serial Stations | 80-0423 |
| 1977-1978 | Current Meter/Wind Records | 80-0423 |
| 1977-1978 | Non-Metal Analysis (Hydrocarbons) | 80-0423 |
| 1977-1978 | Bacteria - Behavior | 80-0423 |
| 1977-1978 | Bacteria - Degradation Rates | 80-0423 |
| 1977-1978 | Bacteria - Enumeration | 80-0423 |
| 1977-1978 | Bacteria - Enumeration | 80-0423 |
| 1977-1978 | Bacteria - Taxonomy/Physiological Diversity | 80-0423 |

| Year | Data Type | NODC Accession Number |
|---|---|---|
| 1977-1978 | Respirometry Experiment | 80-0423 |
| 1977-1978 | Trace Metals - Sediment (Diver Core) | 80-0423 |
| 1977-1978 | Sediment Size Analysis | 80-0423 |
| 1977-1978 | Stomach Contents | 80-0423 |
| 1977-1978 | Demersal Fish | 80-0423 |
| 1977-1978 | Shrimp Bioassay | 80-0423 |
| 1977-1978 | Trace Metals | 80-0423 |
| 1977-1978 | Trapped Suspended Sediment | 80-0423 |
| 1977-1978 | Non-Metal Analysis | 80-0461 |
| 1977-1978 | Bacteria - Behavior | 80-0461 |
| 1977-1978 | Bacteria - Degradation Rates | 80-0461 |
| 1977-1978 | Bacteria - Enumeration | 80-0461 |
| 1977-1978 | Bacteria - Taxonomy/Physiological Diversity | 80-0461 |
| 1977-1978 | Respirometry Experiment | 80-0461 |
| 1977-1978 | Trace Metals - Sediment (Diver Core) | 80-0461 |
| 1977-1978 | Sediment - Size Analysis | 80-0461 |
| 1977-1978 | Stomach Contents | 80-0461 |
| 1977-1978 | Demersal Fish | 80-0461 |
| 1977-1978 | Shrimp Bioassay | 80-0461 |
| 1977-1978 | Trace Metals | 80-0461 |
| 1977-1978 | Trapped Suspended Sediment | 80-0461 |

| Year | Data Type | NODC Accession Number |
|---|---|---|
| 1978-1979 | Stomach Contents | 80-0416 |
| 1978-1979 | Clay Mineralogy | 80-0416 |
| 1978-1979 | Bioassay (Toxicity) | 80-0416 |
| 1978-1979 | Algae | 80-0416 |
| 1978-1979 | Tagging | 80-0416 |
| 1978-1979 | Histopathology and Bacteriology | 80-0416 |
| 1978-1979 | Morphometric | 80-0416 |
| 1978-1979 | Benny Census | 80-0416 |
| 1978-1979 | Biomass Samples - Weight and Barnacles | 80-0416 |
| 1978-1979 | Pistol Shrimp and Stone Crab | 80-0416 |
| 1978-1979 | Biomass - Large Cryptic Samples | 80-0416 |
| 1978-1979 | Surficial Sediments | 80-0416 |
| 1978-1979 | Suspended Particulates | 80-0416 |
| 1978-1979 | Sediments | 80-0416 |

| Year | Data Type | NODC Accession Number |
|---|---|---|
| 1978-1979 | Pb - 210 | 80-0416 |
| 1978-1979 | Bacteria - Enumeration | 80-0416 |
| 1978-1979 | Bacteria - Degradation Rates | 80-0416 |
| 1978-1979 | Bacteria - Taxonomy | 80-0416 |
| 1978-1979 | Bacteria - Growth Characteristics | 80-0416 |
| 1978-1979 | Trace Metals | 80-0416 |
| 1978-1979 | Trace Metals - Organism, Sediment, Water | 80-0416 |
| 1978-1979 | Hydrography | 80-0416 |
| 1978-1979 | Electromagnetic Current Meter | 80-0416 |
| 1978-1979 | Total Suspended Solids | 80-0416 |
| 1978-1979 | Continuous Current Meter | 80-0416 |
| 1978-1979 | Meteorological Data | 80-0416 |
| 1978-1979 | Wave Data | 80-0416 |
| 1978-1979 | Hydrocabons, Biocides and Sulfur | 80-0416 |
| 1978-1979 | Respirometry | 80-0416 |

| Year | Data Type | NODC Accession Number |
|---|---|---|
| 1979-1980 | Data being archived, will be available in late 1980 | TBA |

# INTRODUCTION

## Location of Study Area

The area selected for study is the operational Buccaneer Gas and Oil Field located approximately 49.6 kilometers (26.8 nautical miles) south southeast of the Galveston Sea Buoy off Galveston, Texas (Figure 1). This field was selected in 1975 as the study area because: (a) the field had been in production for about 15 years, which time had allowed full development of the associated marine communities; (b) it was isolated from other fields which facilitated the selection of an unaltered area (for comparison) within a reasonable distance of the field; (c) it produced both gas and oil that represented sources of pollutants from marine petroleum extraction; (d) its location simplified logistics and reduced the cost of the research; and (e) the Texas offshore area had not been fully developed for gas and oil production but was expected to experience accelerated exploitation in the future.

## Operation History of Buccaneer Field

Buccaneer Field was developed by Shell Oil Company in four offshore blocks leased in 1960 and 1968 as follows:

| Year | Lease Number | Block Number | Acreage | Hectares |
|---|---|---|---|---|
| 1960 | G0709 | 288 | 2,790 | 1,129 |
| 1960 | G0713 | 295 | 4,770 | 1,930 |
| 1960 | G0714 | 296 | 4,501 | 1,821 |
| 1968 | G1783 | 289 | 2,610 | 1,056 |

In development of the field, 17 structures were built; two are production platforms, two are quarters platforms, and 13 are satellite structures surrounding well jackets. Initial exploratory drilling began about mid-summer of 1960 with mobile drilling rigs. When (as the result of the exploratory drilling) proper locations for platforms were selected, the permanent production platforms were constructed.

There have been no reports of major oil spills from this field. There have been some reported losses of oil due to occasional mechanical failure of various pieces of equipment. The largest reported spill was three barrels in 1973. The reported oil spill chronology and quantity for Buccaneer Field is as follows:




FIGURE 1. LOCATION OF BUCCANEER FIELD

| Date | Source | Amount | |
|---|---|---|---|
| | | Barrels | Liters |
| September 1973 | Platform 296-B | 0.5 | 79 |
| November 1973 | Unknown | 3.0 | 477 |
| July 1974 | Platform 296-B | 0.5 | 79 |
| August 1974 | Platform 296-B | 1.7 | 265 |
| September 1975 | Platform 288-A | 0.2-0.4 | 38-56 |
| Totals | | 5.9-6.1 | 938-956 |

Buccaneer Field first began operations with the production of oil. Later, when significant quantities of gas were found, the field began producing both oil and gas and has continued to do so to date.

The production platforms and satellites (well jackets) are connected by a number of pipelines with a 50.8 centimeters (20-inch) diameter main pipeline connecting the field to shore. All of the pipelines that are 25.4 centimeters (10 inches) or greater in diameter are buried. The Blue Dolphin Pipeline Company was granted a pipeline permit (No. G1381, Blocks 288 and 296) in 1965 and has operated the pipeline since its construction.

Buccaneer Field occupies a limited area (about 59.3 $km^2$; 22.9 sq. statute miles) leased in the northwestern Gulf of Mexico. Four types of structures are located in Buccaneer Field: production platforms, quarters platforms, satellites (well jackets), and flare stacks. These are shown in Figure 2, which is an oblique aerial photograph of production platform 288-A and vicinity within Buccaneer Field. A map of Buccaneer Field, (Figure 3) depicts the locations of platforms and satellites within the field.




FIGURE 2. BUCCANEER FIELD STRUCTURES




FIGURE 3. SHELL OIL COMPANY'S ALPHANUMERICAL IDENTIFICATION OF BUCCANEER GAS AND OIL FIELD STRUCTURES

# WORK UNIT 2.4.2 - TRACE METALS

Southwest Research Institute

J. B. Tillery

## ABSTRACT

This report provides an overview of four years (1976-1980) of trace metal investigations conducted as a part of a larger multidisciplinary, marine, environmental assessment program of the Buccaneer gas and oil field. Concentrations of Ag, Al, As, Ba, Be, Cd, Co, Cr, Cu, Hg, Fe, Mn, Ni, Pb, Sb, Se, Sr, Tl and Zn have been determined in surficial sediments, subsurface sediments, suspended particulate matter, seawater, produced brine, crude oil and various tissues of biological organisms collected seasonally near two petroleum production platforms. Concentrations of Ba, Cd, Cr, Co, Cu, Mn, Pb, Sr, Hg and Zn in surficial sediments have been related to the platforms or activities on them. Trace metal concentrations in suspended particulate matter are higher than in bottom sediments. Seawater trace metal concentrations are within the range reported for shelf waters. Produced brine discharge have concentrations of Ba, Cd, Cr, Fe, Hg, Mn, Sr, Tl and Zn that are higher than seawater and vary with time. No evidence of excessive bioaccumulation of trace metals in marine organisms from the area around the production platforms was established. Seasonal variations of trace metals were observed in various marine organisms.

CONTENTS


| | Page |
|---|---|
| Abstract | ii |
| Figures | iv |
| Tables | v |
| 1. Introduction | 1 |
| 2. Methods and Materials | 4 |
| 3. Results and Discussion | 11 |
| Sediments | 11 |
| Suspended particulate matter | 20 |
| Seawater and produced brine | 23 |
| Biofouling | 23 |
| Dominant fauna | 23 |
| Crude oil | 32 |
| 4. Conclusions | 34 |
| References | 36 |
| Appendix | |
| A. Analytical methodologies for the fourth year BGOF study | 39 |




XXVIII

FIGURES


| Number | | Page |
|---|---|---|
| 1. | Sampling diagram for sediments. Second year (1977-1978) | 8 |
| 2. | Sampling grid for third year (1978-1979) surficial sediments at platforms 288-A and 296-B | 9 |
| 3. | Sampling grid for fourth year (1979-1980) surficial sediments at platforms 288-A and 296-B | 10 |
| 4. | Scatter plot of surficial sediments Ni concentration versus Fe concentration from 5N $HNO_3$ leach (winter and spring samplings 1978-1979) | 14 |
| 5. | Scatter plot of surficial sediments Cd concentration versus Fe concentration from 5N $HNO_3$ leach (summer and fall samplings 1978-1979) | 15 |
| 6. | Scatter plot of surficial sediments Pb concentration versus Fe concentration from 5N $HNO_3$ leach (summer and fall samplings 1978-1979) | 16 |
| 7. | Scatter plot of surficial sediment Zn concentration versus Fe concentration from 5N $HNO_3$ leach (summer and fall samplings 1978-1979) | 17 |

TABLES


| Number | | Page |
|---|---|---|
| 1. | Estimate of Analytical Error for Third and Fourth Years Sediment Samples | 5 |
| 2. | Estimate of Sampling Variability for Third Year Sediment Samples. Reported as Average Percent Coefficient of Variation | 6 |
| 3. | Estimate of Analytical Error for Third and Fourth Years Biological Samples | 7 |
| 4. | Seasonal Comparison of Average Trace Metal Concentrations (µg/g dry wt) in Surficial Sediments from Platform 296-B | 13 |
| 5. | Significant (95% Level) Accumulations of Trace Metals (µg/g dry wt) in Surficial Sediments at Platform 288-A | 21 |
| 6. | 1979-1980 Seasonal Comparison of Trace Metal Concentrations (µg/g dry wt) in Suspended Particulate Matter from Mid-Depth and Bottom Samples Taken at Platform 296-B | 22 |
| 7. | Mean Trace Metal Concentrations (µg/L) Seawater (Unfiltered) and Produced Brine from Platform 296-B During 1979-1980 | 24 |
| 8. | Mean Trace Metal Concentration (µg/mL) in Produced Brine Samples | 25 |
| 9. | Comparison of Seasonal Variations in Average Trace Metal Concentrations (µg/g dry wt) in Chaetodipterus faber (Spadefish) Muscle Tissue with Other Gulf Studies | 28 |
| 10. | Seasonal Comparison of Chaetodipterus faber (Spadefish) Liver Trace Metal Concentrations (µg/g dry wt) with L. campechanus Liver Tissue from STOCS | 29 |
| 11. | Comparison of Seasonal Variations in Average Trace Metal Concentrations (µg/g dry wt) in Archosargus probatocephalus (Sheepshead) Muscle Tissue with Other Gulf Studies | 30 |
| 12. | Comparison of Average Trace Metal Concentration in Stenotomus caprinus (Longspine Porgy) Muscle Tissue (µg/g dry wt) with Other Gulf of Mexico Studies | 31 |
| 13. | Trace Metal Concentrations (µg/g) in Buccaneer Crude Oil | 33 |




XXX

# SECTION 1

## INTRODUCTION

With rising costs and diminishing supplies of petroleum, the next decade will see an increase in the development of offshore and nearshore petroleum resources. The impact this will have on the marine environment and ecology can be minimized with adequate planning and realistic regulations (1). However, the formulation of these plans and regulations will require a sound, technical data base.

This milestone report is an overview of the four years (1976-1980) of trace metal investigations (Work Group 2.4.2) performed on the "Environmental Assessment of the Buccaneer Gas and Oilfield in the Northwestern Gulf of Mexico" (BGOF).

Funding for this program has come from the Environmental Protection Agency (EPA) and the National Oceanic and Atmospheric Administration (NOAA). Management responsibilities are with the Environmental Research Division of the National Marine Fisheries Service, Southeast Fisheries Center, Galveston Laboratory, Galveston, Texas.

This study has a unique place in relation to the other studies funded by the Federal government that deal with the environmental effects of offshore petroleum production. The Buccaneer Gas and Oilfield study was designed to provide data on the chronic, low-level pollution associated with active petroleum production platforms located in the offshore environment. Results from this (BGOF) program along with the MAFLA Rig Monitoring study (2) and the Louisiana Platform study (3), augmented by data from the BLM-STOCS study (4), the MAFLA Baseline study (5) and the OEI studies (6a,6b) should provide a relatively strong data base from which intelligent plans and regulations can be formulated.

The overall objectives of the Trace Metal Work Groups (2.4.2) were to 1) identify the types and extent of environmental and ecosystems alterations associated with development and release of specific trace metal contaminants from an active gas and oilfield, 2) determine the spatial and temporal distribution of 19 trace metals in the Buccaneer gas and oilfield and the effects they have on the various components of the marine ecosystems, and 3) provide data on trace metal concentrations in various components of the marine ecosystem such that capabilities to describe and predict sources, fate and effects of gas and oilfield trace metal contaminants can be developed.

The BGOF is located approximately 49.6 km south/southeast from the Galveston Sea Buoy off Galveston, Texas. The development and operational history of the BGOF are given elsewhere (7).

The first (1976-1977) and second (1977-1978) years of trace metal investigations of the BGOF were performed by Drs. J. B. Anderson (Rice University, Houston, Texas) and R. R. Schwarzer (Texas Southern University, Houston, Texas).

Drs. Anderson and Schwarzer studied 12 trace metals (Ba, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Sr, and Zn) in 1) surficial sediments, suspended sediments and downcore sediments from a) the BGOF study area, b) near petroleum production platform 288-A, c) satellite structure 288-3, and d) from Galveston Bay to the BGOF at 8-km intervals; 2) produced brine from platforms 288-A and 296-B; 3) fouling organisms and barnacles from the legs of platform 288-A and satellite structure 288-3; and 4) various marine organisms from the vicinity of platform 288-A. Along with these trace metal studies, they also did geophysical studies of the bottom and sediment texture analyses.

The objectives of these first two years of study were to 1) characterize bottom sediments throughout the BGOF, 2) determine if trace metal pollution is present in the BGOF, 3) examine possible sources of metal pollution within the BGOF, and 4) determine the concentrations of trace metals in marine organisms that inhabit the BGOF so estimates of bioaccumulation could be made.

Trace metal investigations during the third (1978-1979) and fourth (1979-1980) years of the BGOF environmental study were performed by the author at Southwest Research Institute, San Antonio, Texas.

The same set of 12 trace metals previously examined were continued in the third year in the following samples: surficial sediments, suspended particulate matter, seawater, produced brine, fouling mat, barnacles, and muscle tissue from selected species of demersal and pelagic fishes and macrocrustaceans.

The objectives of the third year were similar to those listed earlier for the previous years, however, the program was designed so a more detailed examination of the temporal and spatial variations of these 12 trace metals could be delineated. Four consecutive seasons (summer, fall, winter and spring) were sampled.

The fourth and final year of the BGOF study saw a change in the suit of trace metals investigated. Eleven of the original 12 trace metals (Co deleted) were retained and augmented by another seven trace metals (Ag, Al, As, Be, Sb, Se, and Tl) which allowed all 13 priority pollutant metals (8) to be examined.

This limited the objectives of the fourth year to those of the previous year since no data base existed for the new metals added. Only two seasons (summer and winter) were sampled during this final year.

The samples analyzed during the fourth year included: surficial sediments, suspended particulate matter, seawater, produced brine, crude oil, and three tissues from three species of dominate fauna that inhabit the area around the platform structures.

## SECTION 2

## METHODS AND MATERIALS

Details of the sampling and analytical methodologies used in the first and second years are given by Anderson and Schwarzer (9a). Sampling and analytical methodologies for the third year are given elsewhere (10). Fourth year sampling and analytical methodologies were those of the third year with some exceptions. Details of these exceptions are provided in Appendix A.

Analytical and sampling variability for the first and second years of the BGOF are provided elsewhere (9b).

Tables 1 and 3 give the analytical error for sediments and biota, respectively, for the third and fourth years. The larger variability for As, Ba, Be, Cr, Hg, Ni, Sb, and Tl are due to the higher variability of the more sensitive graphite furnace AAS technique and the low concentrations present.

Table 2 is an estimate of the sampling variability expressed as the average of the coefficient of variation ($1\ SD \times 100/\bar{x}$) from two sets of duplicate surficial sediments collected during the summer and winter of the third year (1978-1979). This would include the analytical error from Table 1.

All biological samples consisted of pooled samples and no estimate of sampling error is available on individual organisms for the third (1978-1979) and fourth (1979-1980) years.

Figures 1, 2, and 3 illustrate the sampling grid used at the indicated platforms during the second, third, and fourth years, respectively.

TABLE 1. ESTIMATE OF ANALYTICAL ERROR FOR THIRD AND FOURTH YEARS SEDIMENT SAMPLES

| Metal | Analytical Error* % |
|---|---|
| Ag | 27 |
| As | 25 |
| Ba | 27 |
| Be | 15 |
| Cd | 24 |
| Cr | 10 |
| Cu | 5 |
| Fe | 18 |
| Hg | 7 |
| Mn | 28 |
| Ni | 5 |
| Pb | 3 |
| Sb | 16 |
| Se | 20 |
| Sr | 4 |
| Tl | 20 |
| Zn | 10 |

* $\frac{SD \times 100}{\bar{x}}$

$n = 7$

TABLE 2. ESTIMATE OF SAMPLING VARIABILITY FOR THIRD YEAR SEDIMENT SAMPLES. REPORTED AS AVERAGE PERCENT COEFFICIENT OF VARIATION*

| Metal | Cruise 1 (Summer) | Cruise 3 (Winter) |
|---|---|---|
| Ba | 96 | 8 |
| Cd | 5 | 9 |
| Cr | 79 | 30 |
| Co | 79 | 33 |
| Cu | 129 | 30 |
| Fe | 67 | 35 |
| Pb | 60 | 104 |
| Mn | 66 | 18 |
| Hg | 132 | 127 |
| Ni | 49 | 72 |
| Sr | 27 | 6 |
| Zn | 7 | 131 |

* Analyses of two sets of duplicate surficial sediment samples taken at same sample station.

TABLE 3. ESTIMATE OF ANALYTICAL ERROR FOR THIRD AND FOURTH YEARS BIOLOGICAL SAMPLES

| Metal | Analytical Error* % |
|---|---|
| Ag | 19 |
| As | 107 |
| Ba | 34 |
| Be | 43 |
| Cd | 15 |
| Cr | 42 |
| Cu | 10 |
| Fe | 10 |
| Hg | 42 |
| Mn | 26 |
| Ni | 31 |
| Pb | 12 |
| Sb | 94 |
| Se | 3 |
| Sr | 7 |
| Tl | 175 |
| Zn | 2 |

* $\frac{SD \times 100}{\bar{X}}$

n = 5




Figure 1. Sampling diagram for sediments. Second year (1977-1978)*

*Anderson and Schwarzer, 1979.




Figure 2. Sampling grid for third year (1978-1979) surficial sediments at platforms 288-A and 296-B.




Figure 3. Sampling grid for fourth year (1979-1980) surficial sediments at platforms 288-A and 296-B.

## SECTION 3

## RESULTS AND DISCUSSION

### SEDIMENTS

Early investigations of trace metals in surficial sediments from the BGOF indicated there were higher concentrations within 180 meters of the platform structures (9b).

Anderson and Schwarzer (9b) determined there were accumulations of Ba, Pb, Sr, Hg and Zn in surficial sediments within 180 meters of platform 288-A and well jacket 288-3. Control sediments used for comparison were subsurface sediments taken at the same locations as the surficial sediments. These subsurface sediments represented ambient trace metal concentrations prior to the development of the BGOF. Surficial sediments from the BLM-STOCS study (11) were also used as controls since they represented a relative pristine area of the Gulf of Mexico.

There were higher concentrations of Ba, Pb, Sr, and Zn near the platform (288-A) than well jacket (288-3). This was attributed to the size of the platform relative to the well jacket and the fact that no discharges were associated with the well jacket (9b).

The earlier trace metal investigations of Anderson and Schwarzer (9a,9b) of the BGOF study did not attempt to integrate the sediment geochemical parameters that could cause natural variations in trace metal concentrations. However, this was done later (12) using Q-mode cluster analysis to group sediments with similar characteristics and compare their trace metal concentrations. This revealed an accumulation of Ba, Sr, Cd, Co and Pb in surficial sediments near platform 288-A when compared to "pristine" sediments from 1) BLM-STOCS, 2) BGOF far-field samples, and 3) BGOF subsurface samples.

Sources of these sediment trace metal near the platform structure were suspected of coming from 1) corrosion of platform structures 2) sacrificial cathodes, 3) activities on the platform structures, 4) produced brine discharges, 5) corrosion of metallic debris on the bottom, and 6) engine exhaust from pleasure boats (Pb) (9b).

Based upon the finding from the first and second years, the third and fourth years of the BGOF study concentrated upon the "near-field" (within 100 meters) effect of the platform structures 288-A and 296-B.

Samples of surficial sediments were collected during four consecutive seasons during the third year and two seasons (summer and winter) during the fourth and final year. Samples were taken from the vicinity of platforms 288-A and 296-B during the third and fourth years and also at well jacket 288-5 and flare stack 296-B during the third year.

Concentration gradients of Ba, Cd, Cr, Cu, Mn, Pb, Sr, and Zn that decreased with distance from platforms 288-A and 296-B were observed in the third year (1978-1979). In the fourth year (1979-1980), similar gradients were observed for these same trace metals plus Ag and Hg. Table 4 illustrates these gradients identified during the fourth year sampling at platform 296-B. Each of the axis (north, east, south and west) consists of three samples with the arithmetic mean given. The "structure" station consist of one sample directly under the produced brine discharge pipe. The arithmetic mean ($\bar{x}$), standard deviation (SD) and coefficient of variation (CV%) are given for all samples collected at the platform.

Since these concentration gradients could be caused by natural geochemical variations in the sediments, a technique to distinguish between the natural variability and pollution events is needed. One such technique that has been used successfully (13,14) is to normalize the trace metal concentrations to the hydrated Fe oxide fraction of the sediments. The Fe content of the sediments is high (percent range) when compared to the other trace metals (ppm range) and any abnormal input of trace metals to the sediments would have minimal effect on the Fe content but would have a large impact on the other trace metals. Also, Fe being a transition metal would behave chemically like the other transition metals.

The input of abnormal amounts of trace metals to the sediments would be identified as an enriched ratio of metal concentration to Fe concentration. This is also easy to recognize visually by making a scatter plot of the metal concentration versus Fe concentration and identifying pollution as increased scatter along the Y-axis.

Figures 4, 5, 6 and 7 are examples of these scatter plots for Ni, Cd, Pb and Zn from the third year. These plots consist of all the sediment samples collected during a season. Included on the plots are the regression equation and $r^2$ value (goodness of fit) for each season. As a point of comparison, the regression equation with a 95% confidence interval of sediment data from the STOCS study (11) are included. The STOCS sediments can be considered as coming from a pristine area and representative of ambient trace metal concentrations. Examination of Figure 4 shows that Ni is within the limits of the STOCS Ni data and therefore at ambient concentration in the BGOF sediments. Figures 5, 6 and 7 show there is considerable scatter along the Y-axis for Cd, Pb and Zn as compared to the STOCS confidence limits ($p=0.05$). This indicates there is an abnormal input of Cd, Pb and Zn to the sediments. Generally, the fartherest points out on the Y-axis (i.e., pollution) are those stations closest to the platform structures.

TABLE 4. SEASONAL COMPARISON OF AVERAGE TRACE METAL CONCENTRATIONS (µg/g dry wt) IN SURFICIAL SEDIMENTS FROM PLATFORM 296-B

| Cruise | Axis | Ag | Ba | Cd | Cr | Cu | Hg | Fe* | Mn | Pb | Sr | Zn |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| I (Summer 1979) | N | 0.04 | 57.2 | 0.50 | 8.6 | 6.6 | 0.13 | 0.50 | 165 | 136 | 2188 | 1183 |
| | S | 0.01 | 33.9 | 0.99 | 3.4 | 4.6 | 0.24 | 0.51 | 178 | 18 | 147 | 108 |
| | E | 0.03 | 110 | 0.64 | 9.3 | 8.5 | 0.10 | 0.46 | 126 | 162 | 759 | 1088 |
| | W | 0.01 | 37.3 | 0.35 | 6.9 | 6.9 | 0.37 | 0.59 | 206 | 66 | 139 | 273 |
| | Structure | 0.06 | 88.4 | 0.55 | 10.3 | 40.5 | 0.27 | 0.68 | 194 | 201 | 627 | 928 |
| | $\bar{x}$ | 0.03 | 62.0 | 0.61 | 7.3 | 9.3 | 0.21 | 0.53 | 171 | 104 | 794 | 683 |
| | SD | 0.03 | 47.7 | 0.73 | 5.3 | 10.1 | 0.23 | 0.16 | 41 | 115 | 1548 | 755 |
| | CV% | 93 | 76 | 119 | 73 | 109 | 107 | 30 | 24 | 111 | 195 | 111 |
| | n = 13 | | | | | | | | | | | |
| II (Winter 1979) | N | 0.02 | 74.4 | 0.47 | 13.5 | 8.1 | 0.28 | 0.43 | 187 | 69 | 2155 | 1423 |
| | S | 0.01 | 23.2 | 0.14 | 3.6 | 5.1 | 0.05 | 0.39 | 187 | 12 | 89 | 173 |
| | E | 0.03 | 63.7 | 0.46 | 8.3 | 11.2 | 0.13 | 0.55 | 203 | 48 | 331 | 1266 |
| | W | 0.02 | 30.5 | 0.63 | 5.3 | 17.6 | 0.05 | 0.39 | 162 | 63 | 121 | 487 |
| | Structure | 0.09 | 118.0 | 1.21 | 47.9 | 28.0 | 0.24 | 12.07 | 215 | 276 | 390 | 2289 |
| | $\bar{x}$ | 0.03 | 53.3 | 0.48 | 10.7 | 11.9 | 0.14 | 0.50 | 187 | 66 | 652 | 949 |
| | SD | 0.02 | 43.1 | 0.56 | 12.7 | 12.4 | 0.19 | 0.25 | 36 | 84 | 1665 | 1280 |
| | CV% | 85 | 81 | 116 | 119 | 104 | 140 | 50 | 19 | 127 | 255 | 135 |
| | n = 13 | | | | | | | | | | | |

* Concentration in percent.




I = ⊡ ; $y = 4.59x + 2.21$, $R^2 = 0.3262$, Summer

II = ⊙ ; $y = 7.26x + 0.47$, $R^2 = 0.8517$, Fall

III = STOCS regression, $y = 6.74x + 0.89$, $R^2 = 0.9196$

IIIa = STOCS regression, confidence limit at 95%

**Figure 4. Scatter plot of surficial sediments NI concentration versus Fe concentration from 5N $HNO_3$ leach (summer and fall samplings 1978-1979).**




I = ⊙ ; y = 1.87x + 0.124, $R^2$ = 0.0367, Summer

II = △ ; y = 0.31x + 0.045, $R^2$ = 0.1614, Fall

III = STOCS regression, y = 0.063x + 0.075, $R^2$ = 0.8195

IIIa = STOCS regression, confidence limit at 95%

**Figure 5. Scatter plot of surficial sediments Cd concentration versus Fe concentration from 5N $HNO_3$ leach (summer and fall samplings 1978-1979).**




I = ⊙ ; y = 163x + 74, $R^2$ = 0.0035, Summer

II = △ ; y = -8.8x + 91, $R^2$ = 0.0007, Fall

III = STOCS regression, y = 2.9x + 0.7, $R^2$ = 0.8139

IIIa = STOCS regression, confidence limit at 95%

**Figure 6. Scatter plot of surficial sediments PB concentration versus Fe concentration from 5N $HNO_3$ leach (summer and fall samplings 1978-1979).**




I = ○ ; y = 2994x — 826, $R^2$ = 0.2454, Summer

II = △ ; y = 2634x — 870, $R^2$ = 0.5596, Fall

III = STOCS regression, y = 30x + 2.5, $R^2$ = 0.9754

IIIa = STOCS regression, confidence limit at 95%

**Figure 7. Scatter plot of surficial sediments ZN concentration versus Fe concentration from 5N $HNO_3$ leach (summer and fall samplings 1978-1979).**

Trace metals tend to be complexed by the organics materials present in sediments (15), and there is a positive correlation between the amount of organic material (as measured by TOC) and the amount of trace metals present. This relationship can be used like the one described above for trace metal versus Fe ratios. Enriched ratios indicate abnormal trace metal inputs.

A third parameter can also be used to show trace metal pollution of sediments similar to those described above. It is known that as the grain size of the sediments get smaller, the trace metal concentrations increase. This is caused by the larger surface area available on the smaller sediment particles which afford more binding sites for the metals. This phenomena not only affects the natural levels of trace metals in sediments but also the abnormal inputs (i.e., pollution). Plotting the metal concentration versus the $\phi$ (phi) value (i.e., particle size) of the sediments can be used to identify trace metal pollution as was illustrated earlier for metal-Fe plots.

The metal versus TOC plot and metal versus $\phi$ plots from the third and fourth years have followed the same trends as those illustrated in Figures 4, 5, 6 and 7 for metals versus Fe plots.

The normalizing of the sediment trace metal concentrations to Fe, TOC and $\phi$ indicate that the concentration gradients of Ba, Cd, Cr, Cu, Mn, Pb, Sr, and Zn cannot be attributed to natural variations and are suspected of originating from the platform structures or activities on the platforms. These platform sources may include those listed earlier by Anderson and Schwarzer (9b).

It is interesting that Gallaway et al. (16) has shown that a significant amount of sport fishing occurs at these platforms which reinforces the theory that pleasure boat exhaust are an important source of Pb to the marine sediments near the platforms.

The distribution of trace metals in sediments around the platforms generally follows the seasonal current patterns (9b,10). Currents, wave action and turbulence around the platforms (17) tend to continually redistribute the bottom sediments away from the platforms (18) which effectively prevents any long-term buildup of contaminated sediments near the platform structures.

Estimations of the amounts of trace metals accumulating in the surficial sediments within 180 meters of platforms 288-A and 296-B were made by using mean sediment metal concentrations normalized to Fe. Correction for ambient levels of trace metals in the surficial sediments was made by normalizing (to Fe) the mean trace metal concentrations in the bottom samples of sediment cores taken in the second year (1977-1978) BGOF study. According to Anderson and Schwarzer (9b), these sediments represent the predevelopment period of the BGOF and should be representative of ambient trace metal concentrations for this area of the Gulf.

The mean metal to Fe ratios of the bottom cores were used to adjust the mean Fe concentrations of the surficial sediments to give an estimated ambient level of the metal:

$$\left(\frac{M}{Fe}\right)_{core} \times Fe_{surf} = M_{surf\text{-}amb}$$

$$M_{surf\text{-}detn} - M_{surf\text{-}amb} = M_{surf\text{-}accum}$$

where $\left(\frac{M}{Fe}\right)_{core}$ = mean $\frac{\text{metal concentration}}{\text{Fe concentration}}$ in the bottom core samples

$Fe_{surf}$ = mean Fe concentration in the surficial sediments

$M_{surf\text{-}amb}$ = mean ambient metal concentration in the surficial sediments

$M_{surf\text{-}detn}$ = mean metal concentration determined in the surficial sediments

$M_{surf\text{-}accum}$ = estimated mean metal concentration accumulating in the surficial sediments

The arithmetic mean concentration of each metal was determined from all sampling points and used along with the standard deviation to calculate a lower confidence limit (LCL) at the 95 percent confidence level. The LCL is calculated as

$$LCL_{0.95} = \bar{x} - t_{0.05\ (n-1)} \frac{s}{\sqrt{n}}$$

where $\bar{x}$ = arithmetic mean

$t_{0.05\ (n-1)}$ = Student's t value with n-1 degrees of freedom at the 0.05 significance level

$s$ = estimated standard deviation

$n$ = sample size.

This value represents the minimum trace mean concentration of that metal in the study area, at the 95-percent confidence level and was used as $M_{surf\text{-}detn}$. A similar calculation was used to determine an upper confidence limit (UCL) on the ambient concentration as

$$UCL_{0.95} = \bar{x} + t_{0.05\ (n-1)} \frac{s}{\sqrt{n}}$$

This represents the maximum mean concentration of ambient mean metal concentration in the surficial sediments and was used for $M_{surf-amb}$. Substituting these into the equation above gives a value which, if positive, implies that a significant accumulation of a particular metal occurred in the surficial sediments.

This exercise was also performed using sediment data from STOCS study (11) as representative of ambient trace metal concentrations in sediments. These results are also given in Table 5.

Table 5 shows similar results when using the BGOF bottom core samples and the STOCS data. There are significant ($p=0.05$) short-term accumulations of Ba, Co, Cr, Cu, and Pb in the sediments near the platform structures. The accumulations of Hg, Mn, Sr and Zn appear to be more persistent with time.

## SUSPENDED PARTICULATE MATTER

Examination of trace metal concentrations in suspended particulate matter (SPM) are important because 1) they provide an important transport medium for trace metals in the marine environment and 2) they are an important food source for filter-feeding organisms. Detritus materials in the SPM can absorb or adsorb large amounts of trace metals from the water column. The organic materials in SPM can also complex trace metal (15).

Eventually the trace metals associated with SPM will be deposited in bottom sediment, but before that occurs the SPM can be moved over great distances by wave and wind action and currents.

Anderson and Schwarzer (9a) showed that SPM from the vicinity of platform 288-A contain larger amounts of detritus and organic material from the platform structure or the biota associated with it. These investigators also found a significant amount of metal flakes from the platform structure contaminating their sediment trap samples. Material in these sediment traps indicated that most of it was large enough that it would be deposited close to the platform structures. However, in those sediment traps closest to the bottom about 10% of the material was fine grain sediments. This material was from the nephloid layer normally present near the bottom (except the winter season). This nephloid layer is suspected of being from resuspended bottom sediments (9a,18).

Variability in trace metal concentration of SPM (9a,10) did not allow the identification of trace metals emanating from the platform structures or the produced brine discharges.

Seasonal and sample variability of SPM trace metals from the fourth year are illustrated in Table 6. Trace metal concentrations in SPM are usually higher than the concentration in the bottom sediments (9a,10).

TABLE 5. SIGNIFICANT (95% LEVEL) ACCUMULATIONS OF TRACE METALS (µg/g dry wt) IN SURFICIAL SEDIMENTS AT PLATFORM 288-A

| Trace Metal | Second Year | Third Year | | | | Fourth Year |
|---|---|---|---|---|---|---|
| | | Summer | Fall | Winter | Spring | Summer |
| BGOF Bottom Core Sediments Representative of Ambient[a] | | | | | | |
| Ba | 236 | 45 | NS[b] | NS | NS | 8 |
| Cd[c] | NS | NS | NS | NS | NS | NS[d] |
| Co | 2 | 1 | NS | NS | 1 | ND |
| Cr | NS | 2 | NS | NS | NS | NS |
| Cu | NS | 7 | NS | 1 | NS | NS |
| Hg[c] | NS | 0.04 | NS | 0.03 | 0.02 | 0.02 |
| Mn | 81 | 155 | 67 | 107 | 93 | 137 |
| Ni | NS | NS | NS | NS | NS | NS |
| Pb | 30 | NS | NS | 45 | NS | 43 |
| Sr | 61 | 298 | NS | 144 | 14 | 393 |
| Zn | 71 | 390 | 31 | 332 | NS | 522 |
| STOCS Surficial Sediments Representative of Ambient[e] | | | | | | |
| Ba | 252 | 55 | NS | NS | NS | 17 |
| Cd | NS | NS | NS | NS | NS | NS |
| Cr | 2.2 | 3 | NS | 1 | NS | 1 |
| Cu | 4.1 | 9 | 2 | 3 | 1 | NS |
| Mn | 47 | 136 | 48 | 86 | 80 | 115 |
| Ni | 2 | 2 | 1 | NS | NS | NS |
| Pb | 32 | NS | NS | 46 | NS | 45 |
| Zn | 84 | 398 | 38 | 340 | NS | 530 |

a. BGOF Second Year (Anderson and Schwarzer, 1979). See text for explanation.
b. NS - not significant.
c. Many < values in concentration data from core samples and surficial sediment from second year.
d. ND - not determined.
e. STOCS 1975 Season II (Martin and Holmes, 1976). See text for explanation.

TABLE 6. 1979-1980 SEASONAL COMPARISON OF TRACE METAL CONCENTRATIONS (µg/g dry wt) IN SUSPENDED PARTICULATE MATTER FROM MID-DEPTH AND BOTTOM SAMPLES TAKEN AT PLATFORM 296-B

| Depth | Parameter | Ag | Ba | Cd | Cr | Cu | Hg | Fe | Mn | Pb | Sr | Zn |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| CRUISE I - Summer 1979 | | | | | | | | | | | | |
| Mid-depth | $\bar{x}$ | 28.2 | 103 | <62.2 | 59 | 106 | 0.56 | 1625 | 510 | 134 | 691 | 666 |
| | SD | 81.3 | 79 | -- | 67 | 143 | 0.04 | 2060 | 423 | 173 | 879 | 911 |
| | CV% | 288 | 76 | -- | 113 | 134 | 6 | 127 | 83 | 129 | 127 | 137 |
| | n = 13 | | | | | | | | | | | |
| Bottom | $\bar{x}$ | 3.7 | 157 | <62.2 | 76 | 96 | 1.25 | 1111 | 884 | 124 | 586 | 669 |
| | SD | 0.7 | 149 | -- | 119 | 115 | 2.07 | 1171 | 1071 | 242 | 872 | 1414 |
| | CV% | 18 | 95 | -- | 156 | 120 | 166 | 105 | 121 | 195 | 149 | 211 |
| | n = 12 | | | | | | | | | | | |
| CRUISE II - Winter 1979 | | | | | | | | | | | | |
| Mid-depth | $\bar{x}$ | 1.2 | 748 | 13.6 | 155 | 74 | 2.17 | 700 | 453 | 34 | 174 | 64 |
| | SD | 2.9 | 1069 | 16.7 | 255 | 60 | 1.20 | 519 | 576 | -- | 144 | 44 |
| | CV% | 252 | 143 | 123 | 145 | 82 | 55 | 74 | 127 | -- | 83 | 69 |
| | n = 12 | | | | | | | | | | | |
| Bottom | $\bar{x}$ | 0.3 | 393 | 7.4 | 126 | 126 | 1.00 | 2152 | 1380 | 43 | 199 | 150 |
| | SD | 0.3 | 390 | 8.6 | 264 | 164 | 0.68 | 1393 | 1524 | 24 | 204 | 298 |
| | CV% | 97 | 99 | 117 | 209 | 131 | 68 | 65 | 118 | 54 | 102 | 199 |
| | n = 12 | | | | | | | | | | | |

This is probably a function of the dominant amount of clay in the SPM and the higher amount of organic material present (18).

## SEAWATER AND PRODUCED BRINE

Like the SPM, seawater is an important transport medium of dissolved trace metals. Produced brine discharged into the marine environment can be a potential hazard to marine life if significant quantities of toxic metals are present (19).

Anderson and Schwarzer (9b) reported only Ba and Sr to be in sufficient concentration in produced brine to effect the marine environment. However, the analytical technique used did not allow an accurate determination of the transition metals present.

Results from the third and fourth years of the BGOF study indicate the produced brine discharge have concentrations of Ba, Cd, Cr, Fe, Hg, Mn, Sr, Tl and Zn significantly greater than the seawater to which it is discharged (10) (see Table 7). This could result in an increase of trace metals in the vicinity of the platforms. There are significant seasonal differences in certain metals concentrations in the produced brine discharge that may be the result of discharging from different wells or contamination from the discharge stand pipe. These seasonal differences are shown in Table 8.

Seawater trace metal concentrations from the area around platforms 288-A and 296-B were within the range of noncontaminated shelf waters (20). Concentrations of trace metals in seawater and produced brine samples are given in Table 7.

## BIOFOULING

Barnacles and other marine organisms that attach themselves to the platform structures are a food source for "grazing" fish (16). Any accumulation of trace metals in the fouling organisms could be passed along to other animals higher in the food web. Also these fouling organism would have the most consistent exposure to trace metals from the platform structures.

No excessive accumulation of trace metal has occurred in the fouling organisms (9b,10). Also there is no evidence that fouling organisms close to the produced brine discharge have accumulated more trace metals that other fouling organisms (10).

## DOMINANT FAUNA

Marine organisms that inhabit the waters near sources of trace metal pollution have been used as "pollution indicators" to quantify the type and extent of trace metal pollution (21). It is preferable to use sessile organisms even if they have to be transposed to the

TABLE 7. MEAN TRACE METAL CONCENTRATIONS (µg/L) SEAWATER (UNFILTERED) AND PRODUCED BRINE FROM PLATFORM 296-B DURING 1979-1980

| Sample | Cruise | | Ag | As | Ba | Be | Cd | Cr | Cu | Fe | Hg |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Seawater | Summer 1979 | $\bar{x}$ | 0.033 | <0.037 | 0.121 | <0.002 | 0.050 | 0.051 | 0.340 | 1.24 | 0.032 |
| | | Sd | 0.028 | - | 0.068 | - | 0.049 | 0.029 | 0.174 | 0.33 | 0.004 |
| | | CV % | 85 | - | 56 | 0 | 98 | 57 | 51 | 27 | 13 |
| | | n = 5 | | | | | | | | | |
| | Winter 1979 | $\bar{x}$ | 0.006 | 0.076 | 0.259 | 0.038 | 0.037 | 0.106 | 1.10 | 5.14 | 0.094 |
| | | Sd | 0.001 | 0.050 | 0.161 | 0.002 | 0.018 | 0.020 | 0.43 | 1.94 | 0.019 |
| | | CV % | 17 | 66 | 62 | 6 | 50 | 19 | 40 | 38 | 21 |
| | | n = 4 | | | | | | | | | |
| Produced Brine | Summer 1979 | $\bar{x}$ | 0.049 | 0.091 | 2.40 | <0.002 | 2.06 | 0.879 | 1.83 | 2901 | 0.076 |
| | | Sd | 0.018 | 0.041 | 1.62 | - | 1.92 | 0.761 | 1.76 | 772 | 0.020 |
| | | CV % | 37 | 45 | 68 | - | 93 | 87 | 97 | 27 | 27 |
| | | n = 2 | | | | | | | | | |
| | Winter 1979 | $\bar{x}$ | 0.007 | 0.551 | 10.4 | 0.025 | 0.160 | 0.970 | 1.41 | 2874 | 0.239 |
| | | Sd | 0.002 | 0.058 | 0.6 | 0.005 | 0.004 | 0.392 | 0.72 | 461 | 0.160 |
| | | CV % | 27 | 11 | 6 | 20 | 3 | 40 | 51 | 16 | 67 |
| | | n = 2 | | | | | | | | | |

| Sample | Cruise | | Mn | Ni | Pb | Sb | Se | Sr | Tl | Zn |
|---|---|---|---|---|---|---|---|---|---|---|
| Seawater | Summer 1979 | $\bar{x}$ | 0.064 | 0.148 | 0.225 | 0.111 | <0.027 | 46.5 | 0.053 | 2.59 |
| | | Sd | 0.045 | 0.080 | 0.140 | 0.019 | - | 5.3 | 0.027 | 1.27 |
| | | CV % | 71 | 54 | 62 | 17 | - | 11 | 51 | 49 |
| | | n = 5 | | | | | | | | |
| | Winter 1979 | $\bar{x}$ | 0.073 | 0.648 | 0.728 | 0.051 | 0.057 | 53.7 | <0.050 | 3.43 |
| | | Sd | 0.032 | 0.192 | 0.239 | 0.035 | 0.044 | 4.6 | - | 1.11 |
| | | CV % | 43 | 30 | 33 | 68 | 77 | 9 | - | 32 |
| | | n = 4 | | | | | | | | |
| Produced Brine | Summer 1979 | $\bar{x}$ | 2.69 | 0.354 | 0.610 | 0.087 | <0.027 | 191 | 0.225 | 93.7 |
| | | Sd | 0.48 | 0.165 | 0.459 | 0.001 | - | 151 | 0.088 | 102 |
| | | CV % | 18 | 47 | 75 | 1.1 | - | 79 | 39 | 109 |
| | | n = 2 | | | | | | | | |
| | Winter 1979 | $\bar{x}$ | 5.90 | 1.10 | 0.959 | 0.034 | <0.027 | 269 | 0.425 | 28.1 |
| | | Sd | 1.49 | 0.11 | 0.210 | 0.001 | - | 3 | 0.047 | 23.8 |
| | | CV % | 25 | 10 | 22 | 2 | - | 1 | 11 | 85 |
| | | n = 2 | | | | | | | | |

TABLE 8. MEAN TRACE METAL CONCENTRATION (µg/mL) IN PRODUCED BRINE SAMPLES

| Year | Cruise | Ba | Fe | Sr |
|---|---|---|---|---|
| Second (1977-78) | Summer | 3.5 | 10.0 | 33.5 |
| Third (1978-79) | Summer | 1548 | 340 | 58 |
| | Fall | 1542 | 203 | 64 |
| | Winter | 1029 | 380 | 67 |
| | Spring | 1056 | 118 | 94 |
| Fourth (1979-80) | Summer | 2 | 2901 | 191 |
| | Winter | 10 | 2874 | 269 |

study area. This is not always possible because of technical problems and cost considerations. Other free-moving organisms can be used as indicators of pollution if their life-cycle is limited to a restricted area. Gallaway et al. (16) have shown that spadefish (Chaetodipterus faber) are "habitat faithful" to a particular platform structure and, therefore, may be used as a pollution indicator.

Various marine biota have been collected and analyzed for trace metals over the four years of the BGOF study (9a,10). Perhaps the most important limiting factors in using these various species of biota as pollution indicators is their availability and their natural habitat range. In regards to availability, spadefish (Chaetodipterus faber), sheepshead (Archosargus probatocephalus), sugar shrimp (Trachypenaeus similis) and stone crabs (Menippe mercenaria) appear to be the better candidates. Each of those organisms have been collected at various times during the BGOF study.

Another limiting feature is how to determine bioaccumulation of trace metals in a free-living population of marine animals. Comparison to a similar population in a pristine area or to a population under controlled laboratory conditions are the only technique available. This immediately limits the available organisms to those with a sufficient data base in the literature.

Anderson and Schwarzer (9b) analyzed tissues from several different species of biota during the first and second years of the BGOF study. However, because of a lack of sensitivity for most metals and instrument problems, much of the muscle tissue (i.e., low concentrations) data is of limited value because of the large numbers of data points below the limit of detection.

Comparisons of the first year's trace metal concentrations of Cd, Cr, Cu, Mn, Pb and Zn in shrimp, fish muscle and fish livers with similar data from the STOCS (22) study was presented by Anderson and Schwarzer (9b). The average concentrations of Cd and Cr in shrimp were lower than the STOCS data; the average concentration of Pb and Zn were higher. The average concentrations of Cr, Cu, and Zn were higher in fish muscle tissue than the STOCS averages; Cd, Mn and Pb were lower than the STOCS data. Concentrations of Cd, Cr, Cu, Mn and Pb in fish liver tissues were higher than the average concentrations of the STOCS data; Zn concentrations were lower. No comparisons of the biota trace metal data from the second year were made.

Spadefish (C. faber), sheepshead (A. probatocephalus) and longspine porgy (S. caprinus) muscle tissues were analyzed during the third year. During the fourth year, spadefish (C. faber), sheepshead (A. probatocephalus) and stone crab (M. mercenaria) were analyzed. Also during the fourth year, three different tissues of spadefish (C. faber) and sheepshead (A. probatocephalus) were analyzed: gills, kidney, and muscle. Gonads were to be analyzed but the fish collected did not have sufficiently

developed gonads to sample. Diseased fish and healthy fish were also analyzed (fourth year) to determine if correlation of trace metal concentrations and occurrence of fin rot disease could be established.

Seasonal trace metal concentrations in spadefish (C. faber) muscle tissue from the third (1978-1979) and fourth (1979-1980) years are presented in Table 9. There are significant (p=0.05) seasonal differences in the trace metal concentrations of Fe and Pb. Comparison of the third year (1978-1979) summer data to the data from the Louisiana Platform Study indicates there are significantly (p=0.05) higher concentrations of Cd, Cr, and Cu from the latter study. Considering the other sampling periods, the differences are not significant (p=0.05). This suggests that to determine if bioaccumulation of trace metals has occurred in spadefish (C. faber), consideration of the seasonal variability and possibly the spatial variability will have to be made. Also, a larger sampling population will be needed.

Table 10 is a comparison of the trace metal concentrations in spadefish (C. faber) liver tissue from three years of the BGOF compared to red snapper (L. campechanus) liver data from the STOCS (22). There is significant (p=0.05) seasonal difference for Cu between the second year (1977-1978) and the summer season of the fourth year (1979-1980). There are no significant (p=0.05) differences between the BGOF data and the STOCS data.

Table 11 is a comparison of the seasonal variability (third year) of trace metals in muscle tissue of sheepshead (A. probatocephalus) with similar data from the Louisiana Platform Study (23). There are significant (p=0.05) differences between the summer Cr, Cu, Fe, and Ni concentrations and the Louisiana Platform concentrations. However, the concentrations of these metals from the other seasons are not significantly different than the Louisiana Platform Study concentrations. This suggests the same precautions given earlier about determining bioaccumulations in spadefish (C. faber) should also apply to sheepshead (A. probatocephalus) muscle tissue.

Comparison of diseased and healthy spadefish (C. faber) during the fourth year (1979-1980) indicated only Ba in gills tissue was significantly (p=0.05) higher when compared to healthy gills. This higher Ba concentration (4.1 µg/g) in the gills may be caused by fine grain sediments trapped in the gill tissue rather than bioaccumulation of Ba by the spadefish. The relationship of the higher Ba in the diseased gill tissues should be investigated further to determine if it is a causative factor of the disease or possibly a symptom of it.

During the summer (1978-1979) sampling, no macrocrustacean samples were available at the platform and longspine porgy (Stenotomus caprinus) was substituted. The results of the trace metal analysis of these fish are presented in Table 12. The concentrations of Cr, Fe, Ni, and Pb were significantly (p=0.05) higher than similar data on longspine

TABLE 9. COMPARISON OF SEASONAL VARIATIONS IN AVERAGE TRACE METAL CONCENTRATIONS (µg/g dry wt) IN CHAETODIPTERUS FABER (SPADEFISH) MUSCLE TISSUE WITH OTHER GULF STUDIES

| | Year | Cruise | Parameter | Cd[a] | Cr[a] | Cu[a] | Fe[a] | Pb[a] |
|---|---|---|---|---|---|---|---|---|
| Buccaneer Oilfield | 1978-1979 | Summer n = 8 | $\bar{x}$ | 0.032 | 0.223 | 0.53 | 8.8 | 0.13 |
| | | | SD | 0.022 | 0.061 | 0.08 | 3.3 | 0.04 |
| | | | RSD% | 69 | 27 | 15 | 38 | 31 |
| | | Fall n = 12 | $\bar{x}$ | 0.158 | 0.966 | 1.72 | 10.8 | 0.30 |
| | | | SD | 0.067 | 0.533 | 0.62 | 6.8 | 0.13 |
| | | | RSD% | 42 | 55 | 36 | 63 | 44 |
| | | Winter n = 18 | $\bar{x}$ | 0.132 | 0.764 | 3.25 | 13.3 | 0.53 |
| | | | SD | 0.048 | 0.825 | 4.87 | 5.5 | 0.38 |
| | | | RSD% | 36 | 108 | 150 | 42 | 72 |
| | | Spring n = 12 | $\bar{x}$ | 0.186 | 1.17 | 1.02 | 22.0 | 0.25 |
| | | | SD | 0.232 | 1.15 | 0.20 | 10.6 | 0.22 |
| | | | RSD% | 125 | 98 | 20 | 48 | 90 |
| | 1979-1980 | Summer n = 5 | $\bar{x}$ | 0.082 | 0.236 | 0.65 | 8 | 0.32 |
| | | | SD | 0.062 | 0.007 | 0.15 | 3 | 0.27 |
| | | | RSD% | 76 | 3 | 23 | 36 | 84 |
| | | Winter n = 5 | $\bar{x}$ | 0.019 | 0.201 | 3.14 | 13 | 0.25 |
| | | | SD | 0.017 | 0.004 | 0.25 | 3 | 0.11 |
| | | | RSD% | 93 | 2 | 8 | 21 | 43 |
| Louisiana Platform Study[b] | 1978 | | Range | <0.011-0.158 | <0.20-3.12 | 0.37-1.83 | 8.6-42.8 | <0.11-0.37 |
| | | | $\bar{x}$ | 0.068 | 0.71 | 1.14 | 18.2 | 0.23 |
| | | | SD | 0.070 | 0.91 | 0.34 | 8.3 | 0.31 |
| | | | RSD% | 102 | 127 | 30 | 45.4 | 133 |

a. Those samples at or below detection limit included as detection limit value.
b. Tillery, 1980.

TABLE 10. SEASONAL COMPARISON OF CHAETODIPTERUS FABER (SPADEFISH) LIVER TRACE METAL CONCENTRATIONS (µg/g dry wt) WITH L. CAMPECHANUS LIVER TISSUE FROM STOCS

| Year | Cruise | Parameter | Ba | Cd | Cr | Cu | Fe | Hg | Mn | Ni | Pb | Sr | Z |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| First (1976-77) | Summer | $\bar{x}$ | NR[1] | 6.4 | 6.6 | 72.8 | NR | NR | 4.8 | NR | 9.1 | NR | 86.8 |
| Second (1977-78) | Summer | $\bar{x}$ | 2.0 | 24.4 | 2.5 | 48.7 | 2198 | 0.2 | 8.7 | 1.9 | 4.9 | 1.8 | 131 |
| | | SD | 1.04 | 19.4 | 4.2 | 43.9 | 829 | 0.12 | 2.4 | 2.98 | 3.5 | 0.8 | 22 |
| | | CV% | 52 | 79 | 167 | 90 | 38 | 60 | 28 | 155 | 71 | 45 | 17 |
| | | n=12 | | | | | | | | | | | |
| Fourth (1979-80) | Summer | $\bar{x}$ | 0.41 | 36.8 | 0.31 | 19.3 | 1055 | 0.151 | 8.54 | 1.48 | 0.64 | 3.55 | 177 |
| | | SD | 0.14 | 33.6 | 0.17 | 4.2 | 455 | 0.075 | 2.44 | 1.49 | 0.53 | 1.15 | 163 |
| | | CV% | 35 | 92 | 56 | 22 | 43 | 50 | 29 | 101 | 83 | 32 | 92 |
| | | n=5 | | | | | | | | | | | |
| | Winter | $\bar{x}$ | 0.37 | 1.42 | 0.19 | 40.0 | 1490 | 0.163 | 3.44 | 0.38 | 0.34 | 4.99 | 68 |
| | | SD | 0.99 | 1.32 | 0.02 | 11.6 | 466 | 0.117 | 1.44 | 0.13 | 0.12 | 8.01 | 32 |
| | | CV% | 24 | 93 | 12 | 29 | 31 | 72 | 33 | 34 | 35 | 161 | 47 |
| | | n=5 | | | | | | | | | | | |
| STOCS[3] (1976-77) | Winter Spring Summer | $\bar{x}$ | NR | 3.70 | 2.20 | 14.7 | NR | NR | 3.40 | NR | 2.80 | NR | 146.4 |
| | | Range | | 0.7-6.10 | 2.0-2.2 | 9.0-15.0 | | | 3.00-4.70 | | 1.0-7.60 | | 100-268 |
| | | n=22 | | | | | | | | | | | |

1. NR = not determined.
2. Presley, B. J. and P. N. Booth, 1979, Vol. III, p. E-3.

TABLE 11. COMPARISON OF SEASONAL VARIATIONS IN AVERAGE TRACE METAL CONCENTRATIONS (µg/g dry wt) IN *ARCHOSARGUS PROBATOCEPHALUS* (SHEEPSHEAD) MUSCLE TISSUE WITH OTHER GULF STUDIES

| | Cruise | Parameter | Cr | Cu | Fe | Ni |
|---|---|---|---|---|---|---|
| Buccaneer Oilfield (1978-1979) | Summer | $\bar{x}$[a] | 0.245[b] | 0.59[b] | 8.5[b] | 0.561[b] |
| | | SD | 0.055 | 0.17 | 2.8 | 0.257 |
| | | RSD% | 22 | 29 | 33 | 46 |
| | | n=8 | | | | |
| | Fall | $\bar{x}$ | 0.890 | 1.51 | 9.6 | 0.625 |
| | | SD | 0.566 | 0.62 | 6.4 | 0.193 |
| | | RSD% | 64 | 41 | 67 | 31 |
| | | n=12 | | | | |
| | Winter | $\bar{x}$ | 0.606 | 1.40 | 12.5 | 0.978 |
| | | SD | 0.483 | 0.68 | 3.8 | 0.529 |
| | | RSD% | 80 | 49 | 30 | 54 |
| | | n=19 | | | | |
| | Spring | $\bar{x}$ | 2.65 | 0.99 | 27.3 | 0.532 |
| | | SD | 2.92 | 0.19 | 16.1 | 0.242 |
| | | RSD% | 110 | 19 | 59 | 45 |
| | | n=12 | | | | |
| Louisiana Platform Study[c] | | Range | <0.20-2.03 | 0.57-2.18 | 12.4-35.7 | <0.38-2.33 |
| | | $\bar{x}$ | 1.04[b] | 1.13[b] | 20.6[b] | 1.17[b] |
| | | RSD% | 70 | 46 | 43 | 54 |

a. Those samples at or below detection limit included as detection limit value.
b. Significant ($p = 0.05$) difference.
c. Tillery, 1979.

TABLE 12. COMPARISON OF AVERAGE TRACE METAL CONCENTRATION IN *STENOTOMUS CAPRINUS* (LONGSPINE PORGY) MUSCLE TISSUE (µg/g dry wt) WITH OTHER GULF OF MEXICO STUDIES

| | Cruise | Parameter | Cr[a] | Fe[a] | Ni[a] | Pb[a] |
|---|---|---|---|---|---|---|
| Buccaneer[b] | Summer 1978 | $\bar{x}$ | 0.58 | 22.3 | 1.63 | 0.14 |
| | | RSD% | 110 | 27 | 37 | 36 |
| | | n=23 | | | | |
| | 1977 | $\bar{x}$ | 0.04 | 4.9 | <0.07 | <0.05 |
| | | RSD% | 25 | 18 | 29 | 20 |
| | | n=11 | | | | |
| STOCS[c] | 1976 | $\bar{x}$ | 0.03 | 4.6 | 0.10 | 0.05 |
| | | RSD% | 33 | 41 | 30 | 80 |
| | | n=9 | | | | |

a. Significant (p = 0.05) difference.
b. *Stenotomus caprinus* substituted when no macrocrustacean samples were available.
c. Presley and Booth, 1979, Table 6.8, p. 6-42.

porgy from the STOCS study (22). However, longspine porgy was only sampled this one season and determination of long-term bioaccumulation of these metals is not possible on this limited data.

Sugar shrimp (T. similis) analyzed in the third year had higher Cd-tissue levels than a similar population from the MAFLA Rig Monitoring Study (2) and the SPR program (13). However, the variability in these Cd data from the BGOF make this high Cd data suspect. The high Cd variability comes from one or two pooled samples that have an unusually high Cd content.

Other investigations have identified higher concentations of Ca in kidneys along with supressed levels of Cu and Mg in gonads in Dover sole (Microstomus pacificus) with fin erosion disease (21). The lack of developed gonad tissues in our samples from the winter season prevented this observation from being made in the BGOF study.

CRUDE OIL

Trace metal content of crude oil produced in the BGOF was determined in the fourth year. Table 13 gives the results of this analysis. The only metal that may impact the marine environment in the event of a major oil spill would be Ni. From this data and the reported volume of oil lost to the marine environment in the BGOF (7), crude oil is an insignificant source of trace metals to the BGOF marine environment.

TABLE 13. TRACE METAL CONCENTRATIONS (μg/g) IN BUCCANEER CRUDE OIL*

| Metal | Concentration |
|---|---|
| Ag | <0.011 |
| Al | <0.07 |
| As | 0.116 |
| Ba | 0.001 |
| Be | <0.0001 |
| Cd | <0.002 |
| Cr | 0.002 |
| Cu | 0.014 |
| Fe | 0.057 |
| Hg | 0.055 |
| Mn | 0.002 |
| Ni | 1.34 |
| Pb | <0.02 |
| Sb | 0.012 |
| Se | 0.0001 |
| Sr | 0.02 |
| Tl | 0.0003 |
| Zn | 0.04 |

* Sample collected from on-shore facility by Work Group 3.7 in Spring 1980.

## SECTION 4

## CONCLUSIONS

1. There are seasonal concentration gradients in surficial sediments of Ag, Ba, Be, Cd, Cr, Co, Cu, Mn, Pb, Sr, Hg and Zn within 180 meters of the platform structures.

2. Various normalizing and statistical techniques to eliminate the natural variability of trace metals in surficial sediments indicates an input of Ba, Cd, Cr, Co, Cu, Mn, Pb, Sr, Hg and Zn from the platform structures, activities on or about the structures or engine exhaust from pleasure/service boats.

3. A comparison with other sediments from the Gulf of Mexico shows there have been short-term accumulations of Ba, Co, Cr, Cu, and Pb in the BGOF sediments near the platforms, and more persistent accumulations of Hg, Mn, Sr and Zn have occurred.

4. Distribution of sediments with trace metal accumulations generally follow the seasonal current patterns near the platform structures.

5. Wave action, currents and turbulence around platform structures tend to disperse bottom sediment and prevents a long-term buildup of contaminated sediments at the platforms except those metals noted in 3.

6. Suspended particulate matter have higher trace metal concentrations than bottom sediments.

7. Variability in trace metal concentrations in suspended particulate matter are too great to identify any anthropogenic inputs related to the BGOF.

8. Trace metal concentrations in seawater are within the range reported for nonpolluted shelf waters (unfiltered).

9. There are seasonal variations in the trace metal concentrations in produced brine.

10. Produced brine discharges have concentrations of Ba, Cd, Cr, Fe, Hg, Mn, Sr, Tl and Zn that are significantly higher than the seawater they are discharged into.

11. Large volumes of produced brine could add a significant amount of trace metals to the marine environment but currents and wave action would tend to dilute and move them away from the platforms.

12. There has been no accumulation of trace metals in biofouling communities from the produced brine discharge.

13. Metal concentrations in fish tissues from the first and second years are much higher than those reported in the BLM-STOCS study and the third and fourth years. Lack of sensitivity and analytical problems in the first and second years limits the usefulness of this data.

14. Comparison of data from spadefish and sheepshead tissue (third and fourth years) with similar data from other areas of the Gulf indicate no bioaccumulation has occurred in these species.

15. Indications of bioaccumulation of Cd in longspine porgy tissue (third year) are questionable because of the high variability and will need further study.

16. Bioaccumulation of trace metals in sugar shrimp cannot be confirmed because seasonal variations are unknown.

17. Comparison of diseased and healthy spadefish trace metal burdens (fourth year) indicates Ba levels in gill tissue of diseased fish are higher. Relation of this finding to the disease state are unknown and will require further laboratory investigations.

18. Trace metal concentrations in spadefish and sheepshead muscle tissue (third year) correlate with the suspended particulate trace metal concentrations, i.e., higher in winter season.

19. Crude oil from the BGOF (fourth year) is low in trace metal content. Only Ni is above 1 ppm.

## REFERENCES


1. Frank, R. A. Federal Plan for Ocean Pollution Research, Development, and Monitoring, Fiscal Years 1979-83. Interagency Committee on Ocean Pollution, Research, Development and Monitoring; Federal Coordinating Council for Science, Engineering and Technology. Published by National Oceanic and Atmospheric Administration, 1979.

2. White, T. T., K. E. Turgon and A. W. Blizzard. Rig Monitoring. In: Baseline Monitoring Studies, Mississippi, Alabama, Florida, Outer Continental Shelf, 1975-1976, J. E. Alexander, Project Manager, Volume VI. Prepared by State University System of Florida, Institue of Oceanography. Bureau of Land Management Contract 08550-CT5-30, 1977.

3. Bedinger, C. A., Project Manager. Ecological Investigations of Petroleum Production Platforms in the Central Gulf of Mexico. Final Report, Bureau of Land Management Contract AA551-CT8-17 (in press).

4. Flint, R. W., Program Manager. Environmental Studies, South Texas Outer Continental Shelf, Biology and Chemistry. University of Texas Marine Science Institute report to Bureau of Land Management, Contract AA550-CT6-17, 1979.

5. Dames and Moore. The Mississippi, Alabama, Florida, Outer Continental Shelf Baseline Environmental Survey, 1977-1978. Final Report, J. E. Alexander, Program Manager. Prepared for the Bureau of Land Management, Contract AA550-CT7-34, 1979.

6a. Montalvo, J. G. and D. V. Brady. Part 1: Toxic Metal Determination in Timbalier Bay Water Samples. In: Offshore Ecology Investigations. Gulf University Research Consortium, 1974.

6b. Montalvo, J. G. and D. V. Brady. Part 2: Toxic Metal Determinations in Offshore Water Samples. In: Offshore Ecology Investigations. Gulf University Research Consortium, 1974.

7. Jackson, W. B., ed. Environmental Assessment of an Active Oil Field in the Northwestern Gulf of Mexico, 1977-78. Volume II: Chemical and Physical Investigations. NOAA Annual Report to EPA, Project Number EPA-IAG-D5-E693-EO. NOAA, NTIS Accession No. PB 80107899, 1979. p. 1.

8. USEPA. Water Program: Guidelines Establishing Test Procedures for the Analysis of Pollutants, Federal Register 41(230):52780-52786, 1976.

9a. Anderson, J. B. and R. R. Schwarzer. Sedimentology and Geochemistry. In: Environmental Assessment of an Active Oil Field in the Northwestern Gulf of Mexico, 1976-1977, W. B. Jackson, ed. NOAA Annual Report to EPA, Project EPA-IAG-D5-E693-EO, NOAA, NTIS Accession No. PB 283890, 1977.

9b. Anderson, J. B. and R. R. Schwarzer. Describe the Fine Sediments and Nepheloid Layer of the Oil Field, Focusing upon Their Relationship to Heavy Metal Adsorption/Determine Levels, Pathways and Bioaccumulation of Heavy Metals in the Marine Ecosystem in Oil Field. In: Environmental Assessment of an Active Oil Field in the Northwestern Gulf of Mexico, 1977-1978, W. B. Jackson, ed. Volume III: Chemical and Physical Investigations. NOAA Annual Report to EPA, Project Number EPA-IAG-D5-E693-EO. NOAA, NTIS Accession No. PB 80107899, 1979. p. i.

10. Tillery, J. B. Trace Metals. Volume III. In: Environmental Assessment of Buccaneer Gas and Oil Field in the Northwestern Gulf of Mexico, 1978-1979, W. B. Jackson and E. P. Wilkins, eds., NOAA Technical Memorandum NMFS-SEFC-42, 93 p. Available from NTIS, Springfield, Virginia, 1980.

11. Martin, E. A. and C. W. Holmes. Surficial Sediments-Seasonal Variability in Trace Metals Content. In: Environmental Studies, South Texas Outer Continental Shelf, Geology. Prepared by the Geological Survey for the Bureau of Land Management, 1976.

12. Wheeler, R. B. Environmental Trace Metal Geochemistry of the Buccaneer Oil and Gas Field. M.S. Thesis, Rice University, Houston, Texas, 1979.

13. Shokes, R. F. Characterization of Baseline Conditions at the West Hackberry Offshore Brine Disposal Site - Texoma Group Strategic Petroleum Reserve Program. Final Report prepared by Science Applications Inc. Department of Energy Contract EL-77-C-01-8788, 1978.

14. Trefry, J. H. and B. J. Presley. Heavy Metal Transport from the Mississippi River to the Gulf of Mexico. In: Marine Pollutant Transfer, H. L. Windom and R. A. Duce, eds., Lexington Books, Lexington, Massachusetts, 1976. p. 39.

15. Yen, T. F. and J.I.S. Tang. Chemical Aspects of Marine Sediments. In: Chemistry of Marine Sediments, T. F. Yen, ed. Ann Arbor Science Publishers, Ann Arbor, Michigan, 1977. p. 2.

16. Gallaway, B. J., M. F. Johnson, R. L. Howard, L. R. Martin and G. S. Boland. A Study of the Effect of Buccaneer Oil Field Structures

and Associated Effluents on Biofouling Communities and Atlantic Spadefish. In: Environmental Assessment of an Active Oil Field in the Northwestern Gulf of Mexico, 1977-1978, W. B. Jackson, ed. Volume II: Data Management and Biological Investigations. NOAA Annual Report to EPA, Project Number EPA-IAG-D5-E693-EO. NOAA, NTIS Accession No. PB 80165970, 1979. p 2.3.8-1.

17. Danek, L. and M. S. Tomlinson. Currents and Hydrography. Volume VI. In: Environmental Assessment of Buccaneer Gas and Oil Field in the Northwestern Gulf of Mexico, 1978-1979, W. B. Jackson and E. P. Wilkins, eds., NOAA Technical Memorandum NMFS-SEFC-40, 33 p. Available from NTIS, Springfield, Virginia, 1980.

18. Brooks, J., E. L. Estes and W. H. Huang. Investigations of Surficial Sediments and Suspended Particulates at Buccaneer Field. Volume II. In: Environmental Assessment of an Active Gas and Oil Field in the Northwestern Gulf of Mexico, 1978-1979, W. B. Jackson and E. P. Wilkins, eds., NOAA Technical Memorandum NMFS-SEFC-36, 216 p. Available from NTIS, Springfield, Virginia, 1980.

19. Frank, P. M. and P. B. Robertson. The Influence of Salinity on Toxicity of Cadmium and Chromium to the Blue Grab, Callinectes sapidus. Bull. Environm. Contam. Toxicol. 21:74-78, 1979.

20. Bryan, G. W. Heavy Metal Contamination in the Sea. In: Marine Pollution, R. Johnston, ed. Academic Press, London, 1976. p. 198.

21. Alexander, G. V., D. R. Young, D. J. McDermott, M. J. Sherwood, A. J. Mearns and O. R. Lunt. Marine Organisms in the Southern California Bight as Indicators of Pollution. In: International Conference on Heavy Metals in the Environment, Symposium Proceedings Volume II, Part 2. Toronto, Ontario, Canada, 1975. p. 555.

22. Presley, B. J. and P. N. Boothe. Trace Metals in Epifauna, Zooplankton and Macronekton. In: Environmental studies, South Texas Outer Continental Shelf, 1977, Volume I, Chapter 6. Texas A and M University. Bureau of Land Management Contract AA 550-CT7-11, 1979.

23. Tillery, J. B., R. E. Thomas, and H. L. Windom. Trace Metals in Ecological Investigations of Petroleum Production Platforms in the Central Gulf of Mexico. Bedinger, C. A., Program Manager, Bureau of Land Management, Contract AA551-CTG8-17 (in preparation).

## APPENDIX A

## ANALYTICAL METHODOLOGIES FOR THE FOURTH YEAR BGOF STUDY

1. Sediments - no change from third year.

2. SPM - no change from third year.

3. Seawater and Produced Brine - same extraction procedure used in third year was used for: Ag, As, Cd, Cr, Cu, Hg, Fe, Mn, Ni, Pb, Sb, Se, Tl and Zn.

   The following elements were determined on the raw seawater and produced brine samples by injecting an aliquot into the graphite tube furnace: Al, Ba, Be and Sr.

4. Crude oil - A 78.6 gram (100 mL) sample was digested with $H_2SO_4$ then ashed in a muffle furnace at 350°C. The resulting ash was solubilized with $HNO_3$ (Suprapur) and quantatitively transferred to a 10-mL volumetric flask. Milli-Q water was used to make the sample to volume. The sample was then analyzed by AAS.

5. Biota - all tissue samples were prepared and analyzed by the low-temperature ashing (AAS) technique used for muscle tissue in the third year.

   Stone crab (Menippe mercenaria) samples consisted only of the soft-tissue parts.